



Trifluralin 5G Herbicide

A selective pre-emergence herbicide for control of certain annual grasses and broadleaf weeds in:

- Container Grown Ornamentals
- Landscape Ornamentals
- Nursery Stock
- Groundcovers
- Established Flowers
- Ornamental Bulbs
- Non-Bearing Fruit and Nut Trees and
- Non-Bearing Vineyards
- Christmas Tree Plantations
- Non-Cropland
- Under Paved Surfaces

Active Ingredient:
 trifluralin: a,a,a-trifluoro-2,6-dinitro-N,N-dipropyl-p-toluidine5%
 Other Ingredients95%
 Total.....100%
 Contains 2 lb active ingredient per 40 lb bag.

KEEP OUT OF REACH OF CHILDREN
CAUTION

If you do not understand the label, find someone to explain it to you in detail.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals.

Personal Protective Equipment (PPE):

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical resistant gloves, such as butyl rubber ≥ 11 mils; or natural rubber ≥ 14 mils; or nitrile rubber ≥ 14 mils
- Shoes plus socks

Follow manufacturer's instructions for cleaning/ maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems, enclosed cabs, in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240 (d) (4-6)), the handler PPE requirements may be reduced or modified as specific in the WPS.

| User Safety Recommendations |
|--|
| <p>Users should:</p> <ul style="list-style-type: none"> • Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. • Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. • Remove PPE immediately after handling product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. |

| FIRST AID |
|---|
| <p>If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.</p> <p>If swallowed: Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.</p> <p>If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call poison control center or doctor for treatment advice.</p> <p>If inhaled: Move person to fresh air. If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably by mouth, if possible. Call a poison control center or doctor for further treatment advice.</p> |
| <p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact Chem-Trec at 1-800-424-9300 for emergency medical treatment information.</p> |

Environmental Hazards

This pesticide is extremely toxic to freshwater marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply in a manner that will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not contaminate water when disposing of equipment washwaters or rinsate.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

| Agricultural Use Requirements: |
|--|
| <p>Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.</p> <p>Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.</p> <p>Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.</p> <p>PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil</p> <p>or water, is:</p> <ul style="list-style-type: none"> • Coveralls • Chemical-resistant gloves • Shoes plus socks |

| Non-Agricultural Use Requirements: |
|---|
| <p>The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for Agricultural Pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.</p> <p>Do not enter or allow others to enter the treated area until dusts have settled. If soil incorporation is required after the application, do not enter or allow others to enter the treated area (except those persons involved in the incorporation) until the incorporation is complete. If the incorporation is accomplished by watering-in, do not enter or allow others to enter the treated area until the surface is dry after the watering-in.</p> |

Information

This product is a preemergence herbicide for control of certain annual grasses and broadleaf weeds in container grown ornamentals, landscape ornamentals, nursery stock, groundcovers, established flowers, ornamental bulbs, non-bearing fruit and nut trees and non-bearing vineyards, Christmas tree plantations, non-cropland, and under paved surfaces. Apply this product before or after planting, but prior to germination of target weeds, or immediately after cultivation. Length of weed control will vary with weed population, potting media or soil conditions, temperature, watering regime, and other factors. Following application, user should monitor and observe level of weed control over time to determine when additional applications may be needed. Repeat applications should not be made sooner than 60 days after a previous application of this product. Do not apply over 240 lb per acre total of this product within a 12-month period.

Use Precautions and Restrictions

Do not apply when wind conditions favor drift of this product from the target area. Do not aerially apply this product.

This product does not control established weeds. Existing weeds should be controlled by cultivation or with postemergence herbicides. Weed residues, prunings and trash should be removed or thoroughly mixed into soil prior to treatment. Soil should be in good condition and free of clods at the time of application. A single rainfall or sprinkler irrigation of 0.5 inches or more, or flood irrigation, is required to activate this product. Optimum weed control is obtained when this product is activated within 3 days of application. If rainfall or irrigation has not occurred within 3 days of application and tillage is possible, this product may be activated using cultivation equipment capable of uniformly mixing the herbicide into the upper 1 to 2 inches of soil. Failure to activate this product within 3 days of application may result in erratic weed control.

Optimum weed control will be obtained when followed by overhead irrigation or rainfall within a few hours after surface application.

Weeds Controlled

The following weeds are controlled when this product is applied at 80lb per acre (1.8lb per 1000sq.ft):

| Common Name | Scientific Name |
|----------------------------|--------------------------------|
| barley, hare | <i>Hordeum leporinum</i> |
| barnyardgrass (watergrass) | <i>Echinochloa crus-galli</i> |
| bluegrass, annual | <i>Poa annua</i> |
| brome | <i>Bromus spp.</i> |
| cheat | <i>Bromus secalinus</i> |
| crabgrass | <i>Digitaria spp.</i> |
| cupgrass, southwestern | <i>Eriochloa gracillis</i> |
| fescue, rattail | <i>Vulpia myuros</i> |
| foxtail | <i>Setaria spp.</i> |
| goosegrass | <i>Eleusine indica</i> |
| jungerice | <i>Echinochloa colonum</i> |
| lovegrass | <i>Eragrostis spp.</i> |
| oat, wild | <i>Avena fatua</i> |
| panicum, fall | <i>Panicum dichotomiflorum</i> |
| sandbur, field | <i>Cenchrus incertus</i> |
| signalgrass | <i>Brachiaria spp.</i> |
| sprangletop, Mexican | <i>Leptochloa univervia</i> |
| stinkgrass | <i>Eragrostis ciliarensis</i> |
| witchgrass | <i>Panicum capillare</i> |

| Annual Broadleaf Weeds | |
|------------------------|----------------------------|
| chickweed, common | <i>Stellaria media</i> |
| goosefoot, nettleleaf | <i>Chenopodium murale</i> |
| knotweed, prostrate | <i>Polygonum aviculare</i> |
| kochia | <i>Kochia scoparia</i> |
| lambsquarters, common | <i>Chenopodium album</i> |
| nettle, stinging | <i>Urtica dioica</i> |
| pigweed | <i>Amaranthus spp.</i> |
| purslane, common | <i>Portulaca oleracea</i> |
| thistle, Russian | <i>Salsola iberica</i> |

The following weeds are partially controlled or suppressed when this product is applied at 80lb per acre (1.8lb per 1000sq.ft):

| Common Name | Scientific Name |
|--------------------------|-----------------------------|
| carpetweed | <i>Mollugo verticillata</i> |
| johnsongrass (from seed) | <i>Sorghum halepense</i> |
| panicum, Texas | <i>Panicum texanum</i> |
| pusley, Florida | <i>Richardia scabra</i> |
| shattercane | <i>Sorghum bicolor</i> |

Application Directions

Apply this product using a drop or rotary-type spreader designed to apply granular herbicides or insecticides. Calibrate application equipment prior to use according to manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly. Do not use spreaders that apply material in narrow concentrated bands. Avoid skips or overlaps as poor weed control or plant injury may occur. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at right angles to the first

Application Techniques

When using a drop-type spreader, a splash board mounted under the hopper will provide more even granule distribution.

A chain fastened to the side of the spreader and allowed to drag on the soil surface can be used to mark the edge of the treated swath and help prevent skips or overlaps.

For treating smaller areas or rows of nursery stock or ornamental beds, a hand held or push-type rotary applicator such as whirlybird or cyclone unit is recommended. For hand held units, walk and turn the crank at a constant rate of speed.

A shaker-type applicator made from a small container with holes punched in the bottom is recommended for small, hard to treat areas. Carefully measure the amount of product needed to avoid over-application.

Uses

Container Grown Ornamentals, Landscape Ornamentals, Nursery Stock, Groundcovers, Established Flowers, Ornamental Bulbs, Non-Bearing Fruit and Nut Trees and Non-Bearing Vineyards, and Christmas Tree Plantations

This product may be used as a preemergence treatment for control of certain annual grasses and broadleaf weeds in container grown ornamentals, landscape ornamentals, nursery stock, groundcovers, established flowers, ornamental bulbs, non-bearing fruit and nut trees and non-bearing vineyards and Christmas tree plantations. Apply this product before or after planting but prior to germination of target weeds, or immediately after cultivation.

Refer to the Information section of this label for use precautions and information on application, application rates and weeds controlled.

Special Use Precautions and Restrictions:

- To avoid possible injury, do not apply this product to:
 - nursery forest or Christmas tree seedling beds, cutting beds, or transplant beds
 - unrooted liners or cuttings that have been planted in pots for the first time
 - pots less than four inches wide
 - groundcovers until they are established and well rooted

Do not apply this product to newly transplanted ornamentals, nursery stock, groundcovers, flowers and non-bearing fruit and nut crops and non-bearing vineyards until soil or potting media has been settled by packing and irrigation or rainfall and no cracks are present or injury may occur.

Do not make preplant applications of this product to areas where gladioli corms less than one inch in diameter will be planted or injury may occur.

Do not apply this product in greenhouse or other enclosed structures.

Do not make repeat applications sooner than 60 days after a previous application of this product.

Do not apply over 240 lb per acre total of this product within a 12-month period.

Users who wish to use this product on plant species not recommended on this label may determine suitability for such uses by making trial application of this product at the listed rate to small number of plants. Prior to using this product on a larger number of plants, the treated plants should be observed for signs of herbicidal injury during 30 to 60 days of normal growing conditions to determine if the treatment is non-injurious to the target plant species. **The user assumes responsibility for any plant damage or other liability resulting from the use of this product on plant species not listed on this label.**

This product may be used on the following established plant species when container grown or field grown:

| Common Name | Scientific Name |
|---------------------------------|-----------------------------------|
| Trees | |
| balsam fir | <i>Abies balsamea</i> |
| white fir | <i>Abies concolor</i> |
| albus-flowering maple | <i>Abutilon hybridum</i> |
| luteus-flowering maple | |
| roseus-flowering maple | |
| tangerine-flowering maple | |
| vesuvius red-flowering maple | |
| flame maple | <i>Acer ginnala</i> |
| Norway maple | <i>Acer platanoides</i> |
| red maple | <i>Acer rubrum</i> |
| red sunset maple | |
| silver maple | <i>Acer saccharinum</i> |
| sugar maple | <i>Acer saccharum</i> |
| queen palm | <i>Areacastrum romanzoffianum</i> |
| river birch | <i>Betula nigra</i> |
| paper birch | <i>Betula papyrifera</i> |
| European white birch | <i>Betula pendula</i> |
| bottle tree | <i>Brachychiton populneus</i> |
| black olive | <i>Bucida buceras</i> |
| Chinese chestnut | <i>Castanea mollissima</i> |
| carob | <i>Ceratonia siliqua</i> |
| redbud | <i>Cercis canadensis</i> |
| filicoides-fernspray cypress | <i>Chamaecyparis obtuse</i> |
| gracilis-slender hinoki cypress | |
| swara false cypress | <i>Chamaecyparis pisifera</i> |
| squarrosa-moss cypress | |
| palm | <i>Chamaedorea cataractarum</i> |
| palm | <i>Chamaedorea costaricana</i> |
| Parlor palm | <i>Chamaedorea elegans</i> |
| cloud nine dogwood | <i>Comus florida</i> |
| flowering dogwood | |
| dogwood, kousa | <i>Comus kousa</i> |
| green hawthorn | <i>Crataegus viridis</i> |
| carrot wood | <i>Cupaniopsis anacardiodes</i> |
| Arizona cypress | <i>Cupressus glabra</i> |
| Russian olive | <i>Elaeagnus angustifolia</i> |
| redgum eucalyptus | <i>Eucalyptus camaldulensis</i> |
| mealy eucalyptus | <i>Eucalyptus cinera</i> |
| silver dollar eucalyptus | |
| coolibah tree | <i>Eucalyptus microtheca</i> |
| red ironbark eucalyptus | <i>Eucalyptus sideroxylon</i> |
| figus | <i>Ficus benjamina</i> |
| mini ficus | |
| white ash | <i>Fraxinus americana</i> |
| shamel ash | <i>Fraxinus udhei</i> |
| ginko-maidenhair tree | <i>Ginko biloba</i> |
| honey locust | <i>Gleditsia triacanthos</i> |
| shademaker honey locust | |
| toyon | <i>Heteromeles arbutiflora</i> |
| Florida anise-tree | <i>Illicium floridanum</i> |

Net Weight: 40 lb (18.1 kg)

EPA REG. NO. 10404-118

EPA EST. NO. 961-PA-01 (L), 8378-IN-1 (K), 82757-OH-001 (M), 82757-FL-001 (S), 82757-MA-002 (H), 89135-AZ-001 (A).

First letter of lot code indicates manufacturing site.

F1560

Manufactured for: **LESCO, Inc.**
 1385 East 36th Street
 Cleveland, OH 44114-4114

702307-PG1



Trifluralin 5G Herbicide

This product may be used on the following established plant species when container grown or field grown:

| Common Name | Scientific Name |
|--------------------------------------|---------------------------------|
| eastern red cedar | <i>Juniperus virginiana</i> |
| Japanese larch | <i>Larix kaempferi</i> |
| American sweet gum | <i>Liquidambar styraciflua</i> |
| tuliptree | <i>Liriodendron tulipifera</i> |
| magnolia | <i>Magnolia spp.</i> |
| crabapple | <i>Malus spp.</i> |
| white mulberry | <i>Morus spp.</i> |
| banana | <i>Musa aluminata</i> |
| blackgum | <i>Nyssa sylvatica</i> |
| sourwood | <i>Oxydendrum arboreum</i> |
| Norway spruce | <i>Picea abies</i> |
| pendula-weeping Norway spruce | |
| repens-spreading Norway spruce | |
| dwarf Alberta spruce | <i>Picea glauca conica</i> |
| white spruce | <i>Picea glauca</i> |
| dwarf globe blue spruce | <i>Picea pungens</i> |
| glauca-Colorado blue spruce | |
| hoopsii-hoop's blue spruce | |
| koster-koster blue spruce | |
| bristlecone pine | <i>Pinus aristata</i> |
| canary island pine | <i>Pinus canariensis</i> |
| shore pine, beach pine | <i>Pinus contorta</i> |
| eldarica pine | <i>Pinus eldarica</i> |
| Bosnian pine | <i>Pinus leucodermis</i> |
| pumillo-shrubby Swiss mountain pine | <i>Pinus mugo</i> |
| Austrian black pine | <i>Pinus nigra</i> |
| monterey pine | <i>Pinus radiata</i> |
| red pine | <i>Pinus resinosa</i> |
| loblolly | <i>Pinus taeda</i> |
| white pine | <i>Pinus strobus</i> |
| columnar Scotch pine | <i>Pinus sylvestris</i> |
| Scotch pine | |
| Japanese black pine | <i>Pinus thunbergiana</i> |
| London planetree | <i>Platanus acerifolia</i> |
| American sycamore | <i>Platanus occidentalis</i> |
| California sycamore | <i>Platanus racemosa</i> |
| podocarpus | <i>Podocarpus spp.</i> |
| cottonwood | <i>Populus deltoides</i> |
| Chilean mesquite | <i>Prosopis chilensis</i> |
| yoshino flowering cherry | <i>Prunus yedoensis</i> |
| douglas fir | <i>Pseudotsuga menziesii</i> |
| scarlet oak | <i>Quercus coccinea</i> |
| bear oak | <i>Quercus ilicifolia</i> |
| pin oak | <i>Quercus palustris</i> |
| willow oak | <i>Quercus phellos</i> |
| red oak | <i>Quercus rubra</i> |
| live oak | <i>Quercus virginiana</i> |
| black locust | <i>Robinia pseudoacacia</i> |
| willow | <i>Salix spp.</i> |
| giant sequoia | <i>Sequoiadendron giganteum</i> |
| mahogany | <i>Swietenia mahogani</i> |
| yellow tab | <i>Tabebuia caraiba</i> |
| bald cypress | <i>Taxodium distichum</i> |
| linden | <i>Tilia spp.</i> |
| eastern hemlock | <i>Tsuga canadensis</i> |
| Chinese elm | <i>Ulmus parvifolia</i> |
| Mexican fan palm | <i>Washington robusta</i> |
| Ornamental Shrubs | |
| edward goucher abelia | <i>Abelia grandiflora</i> |
| glossy abelia | |
| abyssinica acacia | <i>Acacia abyssinica</i> |
| prostrate acacia | <i>Acacia redolens</i> |
| shoestring acacia | <i>Acacia stenophylla</i> |
| copper leaf | <i>Acalypha wilkesiana</i> |
| amur maple | <i>Acer ginnala</i> |
| coral bark Japanese maple | <i>Acer palmatum</i> |
| dwarf Japanese maple | |
| century plant | <i>Agave americana</i> |
| false spiraea | <i>Astilbe chinensis</i> |
| Japanese painted fern | <i>Athyrium nipponicum</i> |
| coyotebrush | <i>Baccharis pilularis</i> |
| william penn barberry | <i>Berberis gladwynensis</i> |
| mentor barberry | <i>Berberis mentorensis</i> |
| atropurea-red leaf Japanese barberry | <i>Berberis thunbergii</i> |
| aurea-golden Japanese barberry | |
| crimson pygmy barberry | |
| rose glow barberry | |
| barbara karst | <i>Bougainvillea spp.</i> |

| | |
|------------------------------------|-----------------------------------|
| California gold | |
| pink pixie | |
| scarlet o'hara | |
| temple fire | |
| Texas dawn | |
| Japanese boxwood | <i>Buxus microphylla japonica</i> |
| Korean boxwood | <i>Buxus microphylla koreana</i> |
| common boxwood | <i>Buxus sempervirens</i> |
| lemon bottlebrush | <i>Callistemon citrinus</i> |
| weeping bottlebrush | <i>Callistemon viminalis</i> |
| spring torch scotch heather | <i>Calluna vulgaris</i> |
| sasanqua camellia | <i>Camellia sasanqua</i> |
| Japanese camellia | <i>Camellia japonica</i> |
| feathery cassia | <i>Cassia artemisioides</i> |
| wild lilac | <i>Ceanothus spp.</i> |
| plum yew | <i>Cephalotaxus drupacae</i> |
| snow-in-summer | <i>Cerastium tomentosum</i> |
| kosteri cypress | <i>Chamaecyparis obtusa spp.</i> |
| nana-dwarf hinoki cypress | |
| torulosa cypress | |
| filifera-thread cypress | <i>Chamaecyparis pisifera</i> |
| areca palm | <i>Chrysalidocarpus lutescens</i> |
| summersweet | <i>Clethra alnifolia</i> |
| sibirica-siberian dogwood | <i>Cornus alba</i> |
| baileyi red osier dogwood | <i>Cornus stolonifera</i> |
| flaviramea-yellowtwig dogwood | |
| royal purple smoke tree | <i>Cotinus coggygria</i> |
| coral beauty smoke tree | <i>Cotinus dammeri</i> |
| eichholz smoke tree | |
| praecox-early cotoneaster | <i>Cotoneaster adpressus</i> |
| cranberry cotoneaster | <i>Cotoneaster apiculatus</i> |
| Pyrenees cotoneaster | <i>Cotoneaster congestus</i> |
| Bearberry cotoneaster | <i>Cotoneaster dammeri</i> |
| Himalayan cotoneaster | <i>Cotoneaster Himalayan</i> |
| rock cotoneaster | <i>Cotoneaster horizontalis</i> |
| zabel cotoneaster | <i>Cotoneaster zabelii</i> |
| sago palm | <i>Cycas revoluta</i> |
| hollandia-warminster broom | <i>Cytisus praecox</i> |
| lena-Scotch broom | <i>Cytisus scoparius</i> |
| fragrant daphne | <i>Daphne odora</i> |
| deutzia | <i>Deutzia spp.</i> |
| hopseed bush | <i>Dodonea viscosa</i> |
| fruitland silver berry | <i>Elaeagnus pungens</i> |
| purple bell heather | <i>Erica cinerea</i> |
| mediterranean pink heather | <i>Erica x darleyensa</i> |
| cornish heather | <i>Erica vagans</i> |
| winged euonymus | <i>Euonymus alatus</i> |
| canadale gold euonymus | <i>Euonymus fortunei</i> |
| emerald n gold euonymus | |
| sunspot euonymus | |
| wintercreeper euonymus | |
| silver king euonymus | <i>Euonymus japonica</i> |
| variegated evergreen euonymus | |
| spreading euonymus | <i>Euonymus kiautschovica</i> |
| pineapple guava | <i>Feljoa sellowiana</i> |
| forsythia | <i>Forsythia spp.</i> |
| august beauty gardenia | <i>Gardenia jasminoides</i> |
| gardenia | |
| radican gardenia | |
| salal/lemon leaf | <i>Gaultheria shallon</i> |
| carolina jessamine | <i>Gelsemium sempervirens</i> |
| woadwaxen | <i>Genista pilosa</i> |
| ross estey-hibiscus | <i>Hibiscus rosa-sinensis</i> |
| rose of sharon-heart | <i>Hibiscus syriacus</i> |
| rose of sharon-red bird | |
| rose of sharon-woodbridge | |
| holly | <i>Ilex spp.</i> |
| mystery gardenia | <i>Illicium annisatum</i> |
| henry garnet holly leaf sweetspire | <i>Itea ilicifolia</i> |
| ixora | <i>Ixora collinea</i> |
| juniper | <i>Juniperus spp.</i> |
| mountain laurel | <i>Kalmia latifolia</i> |
| crepe myrtle | <i>Lagerstroemia indica</i> |
| lantana | <i>Lantana spp.</i> |
| coast leucothoe | <i>Leucothoe axillaris</i> |
| drooping leucothoe | <i>Leucothoe fontainesiana</i> |
| privet | <i>Ligustrum spp.</i> |
| Chinese fountain palm | <i>Livistona chinensis</i> |
| flowering woodbine | <i>Lonicera periclymenum</i> |
| serotina woodbine | |
| trumpet honeysuckle | <i>Lonicera sempervirens</i> |
| leather leaf mahonia | <i>Mahonia bealei</i> |
| creeping mahonia | <i>Mahonia repens</i> |
| wax myrtle | <i>Myrica cerifera</i> |
| compacta-dwarf heavenly bamboo | <i>Nandina domestica</i> |
| harbour dwarf-heavenly bamboo | |
| heavenly bamboo | |
| nana-compacta-heavenly bamboo | |
| nana-purpurea-heavenly bamboo | |
| woods dwarf heavenly bamboo | |
| hardy red oleander | <i>Nerium oleander</i> |
| oleander | |
| ruby lace oleander | |
| fortunes osmanthus | <i>Osmanthus fortunei</i> |
| mockorange | <i>Philadelphus spp.</i> |

| | |
|-------------------------------------|------------------------------------|
| pygmy date palm | <i>Phoenix roeloenii</i> |
| fraser's photinia | <i>Photinia fraseri</i> |
| Japanese andromeda | <i>Pieris japonica</i> |
| mountain fire lily-of-the-valley | |
| snowdrift lily-of-the-valley | |
| templebells lily-of-the-valley | |
| valley rose lily-of-the-valley | |
| valley valentine lily-of-the-valley | |
| forest flame lily-of-the-valley | <i>Pieris japonica x forestii</i> |
| mugo-mugo pine | <i>Pinus mugo</i> |
| green pittosporum | <i>Pittosporum tobira</i> |
| Japanese pittosporum | |
| wheeler's dwarf pittosporum | |
| blue cape plumbago | <i>Plumbago ariculata</i> |
| plumbago | <i>Plumbago capensis</i> |
| yewpine | <i>Podocarpus macrophyllus</i> |
| tassel fern | <i>Polystichum polyblepharum</i> |
| cinquefoil | <i>Potentilla spp.</i> |
| carolina laurel cherry | <i>Prunus caroliniana</i> |
| dwarf pink flowering almond | <i>Prunus glandulosa</i> |
| pyracantha | <i>Pyracantha spp.</i> |
| charisma-monruce raphiolepis | <i>Raphiolepis indica</i> |
| enchantress-mones raphiolepis | |
| India hawthorn | |
| springtime-monme raphiolepis | |
| roundleaf raphiolepis | <i>Raphiolepis ovata</i> |
| azalea/rhododendron | <i>Rhododendron spp.</i> |
| Africa sumac | <i>Rhus lancea</i> |
| ramanas rose | <i>Rosa rugosa</i> |
| rosemary | <i>Rosmarinus officinalis</i> |
| Japanese skimmia | <i>Skimmia japonica</i> |
| reeve's skimmia | <i>Skimmia revesiana</i> |
| Anthony waterer spiraea | <i>Spiraea bumalda</i> |
| dolchia spiraea | <i>Spiraea japonica</i> |
| Japanese alpine spiraea | |
| shirobana spiraea | |
| bridal wreath | <i>Spiraea vanhouttii</i> |
| Chinese lilac | <i>Syringa rothomagensis</i> |
| common lilac | <i>Syringa vulgaris</i> |
| upright Japanese yew | <i>Taxus cuspidata</i> |
| anglojap yew | <i>Taxus media</i> |
| cape honeysuckle | <i>Tecomaria capensis</i> |
| | <i>Temstroemia gymnanthera</i> |
| American arborvitae | <i>Thuja occidentalis</i> |
| emerald arborvitae | |
| globosa-globe arborvitae | |
| little giant-dwarf arborvitae | |
| nigra-dark American arborvitae | |
| pyramidalis-pyramid arborvitae | |
| rheingold arborvitae | |
| techny arborvitae | |
| aureus nana-dwarf golden arborvitae | <i>Thuja orientalis</i> |
| minima glauca-dwarf arborvitae | |
| Christmas palm | <i>Veitchia merrilli</i> |
| viburnum | <i>Viburnum spp.</i> |
| weigela | <i>Weigela spp.</i> |
| xylosma | <i>Xylosma congestum</i> |
| yucca | <i>Yucca filamentosa</i> |
| Groundcovers | |
| wooly yarrow | <i>Achillea tomentosa</i> |
| lily-of-the-nile | <i>Agapanthus spp.</i> |
| carpet bugle | <i>Ajuga spp.</i> |
| beechgrass | <i>Ammophila breviligulata</i> |
| cape weed | <i>Arctotheca calendula</i> |
| thrift | <i>Armeria maritima</i> |
| asparagus fern | <i>Asparagus densiflorus</i> |
| bellflower | <i>Campanula spp.</i> |
| variegated carex | <i>Carex spp.</i> |
| largeleaf iceplant | <i>Carpobrotus edulis</i> |
| dwarf plumbago | <i>Ceratostigma plumbaginoides</i> |
| rockrose | <i>Cistus spp.</i> |
| coreopsis | <i>Coreopsis spp.</i> |
| crown vetch | <i>Coronilla vana</i> |
| pampas grass | <i>Cortadenia selloana</i> |
| cotoneaster | <i>Cotoneaster spp.</i> |
| white iceplant | <i>Delosperma alba</i> |
| descampsia | <i>Descampsia caespitosa</i> |
| trailing rosea iceplant | <i>Drosanthemum floribundum</i> |
| iceplant | <i>Drosanthemum hispidum</i> |
| blue fescue | <i>Festuca ovina glauca</i> |
| strawberry, beach | <i>Fragaria chiloensis</i> |
| gazania | <i>Gazania spp.</i> |
| golden hakonechloa | <i>Hakonechloa macroaureola</i> |
| Algerian ivy | <i>Hedera canariensis</i> |
| English ivy | <i>Hedera helix</i> |
| daylily | <i>Hemerocallis spp.</i> |
| rupturewort | <i>Herniaria glabra</i> |
| albo-marginata hosta | <i>Hosta lancifolia</i> |
| aaronsbeard | <i>Hypericum spp.</i> |

| | |
|----------------------------|-----------------------------------|
| St. Johnswort | |
| angelwing jasmine | <i>Jasminum nitidum</i> |
| trailing iceplant | <i>Lampranthus spectabilis</i> |
| white lily turf | <i>Liriope gigantea</i> |
| big blue ivy | <i>Liriope muscari</i> |
| lilac beauty lily turf | |
| majestic lily turf | |
| silvery sunproof lily turf | |
| variegated liriope turf | |
| green/creeping lily turf | <i>Liriope spicata</i> |
| eulalia grass | <i>Miscanthus sinensis</i> |
| creeping wirevine | <i>Muehlenbeckia axillaris</i> |
| myoporum | <i>Myoporum laetum</i> |
| dwarf mondo grass | <i>Ophiopogon japonicus</i> |
| mondo grass | |
| trailing African daisy | <i>Osteospermum fruticosum</i> |
| Japanese spurge | <i>Pachysandra terminalis</i> |
| fountain grass | <i>Pennisetum alopecuroides</i> |
| ribbon grass | <i>Phalaris arundinacea picta</i> |
| stonecrop (sedum) | <i>Sedum spp.</i> |
| germander | <i>Teucrium chamaedrys</i> |
| Asian jasmine | <i>Trachelospermum asiaticum</i> |
| verbena | <i>Verbena spp.</i> |
| speewell | <i>Veronica spp.</i> |
| periwinkle | <i>Vinca spp.</i> |
| Established Flowers | |
| yarrow | <i>Achillea spp.</i> |
| floss flower | <i>Ageratum houstonianum</i> |
| alysium | <i>Alyssum spp.</i> |
| snapdragon | <i>Antirrhinum majus</i> |
| African daisy | <i>Arctotis spp.</i> |
| silver mound | <i>Artemisia schmidtiana</i> |
| dusty miller | <i>Artemisia stellerana</i> |
| aster (perennial) | <i>Aster spp.</i> |
| begonia | <i>Begonia spp.</i> |
| calendula | <i>Calendula officinalis</i> |
| cornflower | <i>Centaurea cyanus</i> |
| velvet centaurea | <i>Centaurea gymnocarpa</i> |
| sweet sultan | <i>Centaurea moschata</i> |
| chrysanthemum | <i>Chrysanthemum spp.</i> |
| morningglogy | <i>Convolvulus spp.</i> |
| coreopsis | <i>Coreopsis spp.</i> |
| cosmos | <i>Cosmos spp.</i> |
| dahlia | <i>Dahlia spp.</i> |
| dianthus | <i>Dianthus spp.</i> |
| bleeding heart | <i>Dicentra eximia</i> |
| Foxglove | <i>Digitalis mertonensis</i> |
| common foxglove | <i>Digitalis purpurea</i> |
| marigold, cape | <i>Dimorphotheca spp.</i> |
| purple coneflower | <i>Echinacea purpurea</i> |
| California poppy | <i>Eschscholzia californica</i> |
| snow-on-the-mountain | <i>Euphorbia marginata</i> |
| geranium | <i>Geranium spp.</i> |
| geum | <i>Geum spp.</i> |
| gaillardia | <i>Gaillardia spp.</i> |
| gladiolus | <i>Gladiolus spp.</i> |
| guara | <i>Guara lindheiveri</i> |
| baby's breath | <i>Gypsophila paniculata</i> |
| sunflower | <i>Helianthus spp.</i> |
| coral bells | <i>Heuchera spp.</i> |
| aaron's beard | <i>Hypericum calycinum</i> |
| balsam | <i>Impatiens balsamina</i> |
| impatiens | <i>Impatiens spp.</i> |
| iris | <i>Iris spp.</i> |
| ixora | <i>Ixora spp.</i> |

Net Weight: 40 lb (18.1 kg)

EPA REG. NO. 10404-118
EPA EST. NO. 961-PA-01 (L), 8378-IN-1 (K), 82757-OH-001 (M), 82757-FL-001 (S), 82757-MA-002 (H), 89135-AZ-001 (A).

First letter of lot code indicates manufacturing site.

F1560

Manufactured for: LESCO, Inc.
1385 East 36th Street
Cleveland, OH 44114-4114

702307-PG2



Trifluralin 5G Herbicide

This product may be used on the following established plant species when container grown or field grown:

| Common Name | Scientific Name |
|--|---------------------------------|
| Established Flowers (continued) | |
| sweet pea | <i>Lathyrus spp.</i> |
| English lavender | <i>Lavandula angustifolia</i> |
| shasta daisy | <i>Leucanthemum superbum</i> |
| gayfeather | <i>Liatris spicata</i> |
| statice | <i>Limonium spp.</i> |
| lobelia | <i>Lobelia spp.</i> |
| sweet alyssum | <i>Lobularia maritima</i> |
| lupine | <i>Lupines spp.</i> |
| stock | <i>Matthiola spp.</i> |
| four o' clock | <i>Mirabilis jalapa</i> |
| bee balm | <i>Monarda didyma</i> |
| forget-me-not | <i>Myosotis spp.</i> |
| nicotiana | <i>Nicotiana spp.</i> |
| ozark sundrop | <i>Oenothera missouriensis</i> |
| poppy, California | <i>Papaver spp.</i> |
| petunia | <i>Petunia hybrida</i> |
| beard tongue | <i>Penstemon barbatus</i> |
| Russian sage | <i>Perovskia atriplicifolia</i> |
| phlox | <i>Phlox spp.</i> |
| portulaca | <i>Portulaca grandiflora</i> |
| rose | <i>Rosa spp.</i> |
| blackeyed susan | <i>Rudbeckia hirta</i> |
| golden glow | <i>Rudbeckia laciniata</i> |
| salvia | <i>Salvia spp.</i> |
| pincushion flower | <i>Scabiosa spp.</i> |
| stonecrop | <i>Sedum spectabile</i> |
| lamb's ears | <i>Stachy's spp.</i> |
| stoke's aster | <i>Stokesia laevis</i> |
| marigold | <i>Tagetes spp.</i> |
| painted daisy | <i>Tanacetum coccineum</i> |
| nasturtium | <i>Tropaeolum spp.</i> |
| speedwell | <i>Veronica spicatum</i> |
| vinca | <i>Vinca spp.</i> |
| zinnia | <i>Zinna spp</i> |

Ornamental Bulbs

This product may be applied for control of susceptible annual weeds in ornamental bulbs, e.g., bulbous iris, daffodil (narcissus), hyacinth and tulip. Apply this product to the soil surface 2 to 4 weeks after planting, but prior to the emergence of annual weeds. This product may also be applied following bulb emergence. For fall planted bulbs, apply this product again in late winter or early spring to weed-free soil surfaces.

Non-Bearing Fruit and Nut Trees and Non-Bearing Vineyards

Non-bearing fruit and nut trees and non-bearing vineyards are defined as plants that will not bear fruit for at least one year after treatment.

| | | |
|---------------|-----------------|-----------------|
| almond | fig | orange |
| apple | filbert | peach |
| apricot | gooseberry | pear |
| avocado | grape, American | pecan |
| blackberry | grape, European | pistachio |
| blueberry | grapefruit | plum |
| boysenberry | kiwi | pomegranate |
| cherry, sour | lemon | prune |
| cherry, sweet | loganberry | raspberry |
| currant | macadamia nut | walnut, black |
| dewberry | nectarine | walnut, English |
| elderberry | olive | |

Christmas Tree Plantations

Apply this product to established plantings of field grown Christmas tree species listed on this label. Do not apply to seedbeds or seedling transplant beds. Apply only to established plantings. Established plants are defined as those that have been transplanted into their final growing location for a sufficient period of time to allow the soil to be firmly settled around the roots from packing and rainfall or irrigation. Follow all instructions provided in the Information section of this label.

Non-Cropland Areas

This product may be used as a preemergence herbicide for control of certain annual grasses and broadleaf weeds on industrial sites, utility substations, highway guard rails, sign posts and delineators.

Apply this product anytime prior to germination of target weeds. Areas to be treated should be free of established weeds or existing weeds should be controlled with postemergence herbicides.

Refer to the Information section prior to using this product on non-cropland.

| Under Paved Surfaces | | | | | | |
|---|--|--|---------------|-----------------------|---------|---------|
| Site Preparation This product should be used only where the soil to be treated has been prepared according to good construction practices. If rhizomes, stolons, tubers or other vegetative plant parts are present in the site, they should be removed by scalping with grader blade to a depth sufficient to ensure their complete removal. | | | | | | |
| Application Application may be made after the final road bed is established or after the base rock has been added. Do not move soil after application of this product and do not apply this product to soil where asphalt may be applied directly on top of the treated soil. Paving should follow applications of this product as soon as possible. Apply this product to ensure thorough coverage of the base rock layer. Apply with any granular applicator that will apply uniformly. | | | | | | |
| Apply the following amount of this product: | | | | | | |
| <table border="1"> <thead> <tr> <th colspan="2">Amount of LESCO Trifluralin 5G Herbicide</th> </tr> <tr> <th>Per Acre (lb)</th> <th>Per 1000 sq. ft. (lb)</th> </tr> </thead> <tbody> <tr> <td>240-320</td> <td>5.5-7.3</td> </tr> </tbody> </table> | Amount of LESCO Trifluralin 5G Herbicide | | Per Acre (lb) | Per 1000 sq. ft. (lb) | 240-320 | 5.5-7.3 |
| Amount of LESCO Trifluralin 5G Herbicide | | | | | | |
| Per Acre (lb) | Per 1000 sq. ft. (lb) | | | | | |
| 240-320 | 5.5-7.3 | | | | | |

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in original container only. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available, or dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully and completely. It is impossible to eliminate all risks inherently associated with the use of this product. To the extent consistent with applicable law, Buyer and/or User assume all risks of ineffectiveness or other unintended consequences or damages that may result from conditions outside or beyond the control of LESCO, Inc. including but not limited to, such factors as manner of use or application, weather or weather conditions outside the range considered normal at the application site or for the time period in which the product is applied, the presence of other materials, incompatible products, or other influencing factors which are beyond the control of LESCO, Inc.. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and/or User, and Buyer and/or User agrees to hold LESCO, Inc. harmless for any claims relating to such factors.

LESCO, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with the Directions for Use under normal use conditions. To the extent consistent with applicable law, this warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of LESCO, Inc. and Buyer and/or User assume the risk of any such use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, LESCO, INC. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE BUYER AND/OR USER AND THE EXCLUSIVE LIABILITY OF LESCO, INC. FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF LESCO, INC. REPLACEMENT OF THE PRODUCT, OR IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL LESCO, inc., BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES. LESCO, Inc. offers this product, and Buyer and/or User accepts it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of LESCO, Inc.

Information concerning the raw materials composing this product can be obtained by writing to: LESCO, Inc., Attn: RA Dept., 1385 East 36th Street, Cleveland, Ohio 44114-4114, referring to the item number found on this bag.

LESCO is registered trademark and the sweeping design is a trademark of LESCO Technologies, LLC. All other trademarks are the property of their respective owners.

(020510) Rev 10/21/16 VT
Net Weight: 40 lb (18.1 kg)

EPA REG. NO. 10404-118
EPA EST. NO. 961-PA-01 (L), 8378-IN-1 (K), 82757-OH-001 (M), 82757-FL-001 (S), 82757-MA-002 (H), 89135-AZ-001 (A).

First letter of lot code indicates manufacturing site.

F1560
Manufactured for: **LESCO, Inc.**
1385 East 36th Street
Cleveland, OH 44114-4114

702307-PG3